

Aquaculture and the Different Perspectives

Dini Chandran C S and Dr Rekha Sivadasan

*The Cashew Export Promotion Council of India,
Kollam, Kerala*

Abstract

Aquaculture is the rearing of freshwater and saltwater populations under controlled or semi-natural conditions. It includes the business of producing aquatic animals and plants in managed, unnatural aquatic ecosystems for profit (Boyd and Schmittou, 1999). According to the Food and Agriculture Organization (FAO), aquaculture is understood to mean the farming of aquatic organisms including fish, mollusks, crustaceans and aquatic plants. Particular kinds of aquaculture include fish farming, shrimp farming, oyster farming, mariculture, alga culture such as seaweed farming and the cultivation of ornamental fish. Since in 1970s, global aquaculture production has grown rapidly and is now among the fastest growing food production sectors in many countries. Although Asia is by far the leading region contributing to about 90% aquaculture production, the importance of aquaculture is still growing and there is vast scope for its development. Despite its potential, aquaculture is an often-neglected option in agricultural development. Worldwide, millions of people suffer from protein malnutrition causing general ill health. (Olaf Muller and Michael Krawinkel., 2005) In India, fish in the diet is most important in the east and south India. Fish is a nutrient dense food. Fish gives sufficient amounts of all the essential amino acids required by the body for growth and maintenance. Fish is low in fat and high in easily digestible protein and Vitamin B-12. Some fishes are high in heart friendly cholesterol-lowering oils. It's a good source of protein-most varieties contain around 23.5% of protein, the same as meat. In 2015, fish accounted for about 17 per cent of animal protein consumed globally, and provided about 3.2 billion people with almost 20 per cent of their average per capita intake of animal protein. (The state of world fisheries and aquaculture., 2020).

Keywords: Aquaculture, Antibiotic resistance, public health, antimicrobials

INTRODUCTION

Almost all fish produced from aquaculture is mainly for human consumption. Diseases are one of the major constraints to aquaculture and the main limiting factor for economic and socio-economic developments in India and as in many other countries of the world. (De Silva, S.S.2001). It is understood that occurrence of disease is a result of the complex interaction between the host, the pathogen and the environment. Different stress factors such as non-optimal water quality, higher microbial load, poor nutritional status, high stocking density can trigger the chances of infection by opportunistic pathogens in aquatic environment. (Mishra et al., 2017). Most bacterial, parasitic and fungal pathogens are not strictly parasitic micro-organisms. These pathogens have a high adaptability to environmental changes. If the conditions for parasitism are unsuitable, saprophytic relationship will occur. Environmental stress factors can result in increased development of fungal infections. (Ana Cao et al., 2013). Most of the farmers are with a little knowledge of aquaculture health management and with inadequate opportunities for the improvement of disease problems. Most of them do not understand the signs of diseases. For this reason, efforts have to give not only on the occurrence of disease and pathogens but also on the awareness developmental programmes on farmer's disease awareness. Some diseases have caused serious damage, not only the livelihood of fish farmers, but also, to the future development of the industry. (Rosamond L Naylor et al., 2021) Many diseases affecting present day aquaculture is due to the resultant of boosting of culture practices without the basic insight of tangled balance between host, pathogen and environment. (MD. Ashihur Rahman et al., 2019). It is understood that occurrence of disease is a result of the complex interaction between the host, the pathogen and the environment. (Ayalew Assefa and Fufa Abunna., 2018) In conclusion, rather than trying to treat every disease case, it advisable to follow a preventive approach before the event of any disease outbreaks. Besides implementation of better management practices it is relevant to prevent frequent occurrence of disease and production loss in aquaculture. It is clear that aquaculture is a huge industry operating worldwide and growing rapidly. Among these challenges, infectious diseases take the lion share causing billion dollar loss annually. Therefore, problems planning impediment and control strategy based on globally accepted principles and locally applicable strategies are to be followed. These strategies should focus on preventing the development of infection rather than treating diseased stocks.

COMMON DISEASES AFFECTING AQUACULTURE

Almost all fish produced from aquaculture is mainly for human consumption. Diseases are one of the major constraints to aquaculture and the main limiting factor for economic and socio-economic developments in India and as in many other countries of the world. (De Silva, S.S.2001). It is understood that occurrence of disease is a result of the complex interaction between the host, the pathogen and the environment. Different stress factors such as non-optimal water quality, higher microbial load, poor nutritional status, high stocking density can trigger the chances of infection by opportunistic pathogens in aquatic environment. (Mishra et al., 2017). Most bacterial, parasitic and

fungus pathogens are not strictly parasitic micro-organisms. These pathogens have a high adaptability to environmental changes. If the conditions for parasitism are unsuitable, saprophytic relationship will occur. Environmental stress factors can result in increased development of fungal infections. (Ana Cao et al., 2013). Most of the farmers are with a little knowledge of aquaculture health management and with inadequate opportunities for the improvement of disease problems. Most of them do not understand the signs of diseases. For this reason, efforts have to give not only on the occurrence of disease and pathogens but also on the awareness developmental programmes on farmer's disease awareness. Some diseases have caused serious damage, not only the livelihood of fish farmers, but also, to the future development of the industry. (Rosamond L Naylor et al., 2021) Many diseases affecting present day aquaculture is due to the resultant of boosting of culture practices without the basic insight of tangled balance between host, pathogen and environment. (MD. Ashihur Rahman et al., 2019). It is understood that occurrence of disease is a result of the complex interaction between the host, the pathogen and the environment. (Ayalew Assefa and Fufa Abunna., 2018) In conclusion, rather than trying to treat every disease case, it advisable to follow a preventive approach before the event of any disease outbreaks. Besides implementation of better management practices it is relevant to prevent frequent occurrence of disease and production loss in aquaculture. It is clear that aquaculture is a huge industry operating worldwide and growing rapidly. Among these challenges, infectious diseases take the lion share causing billion dollar loss annually. Therefore, problems planning impediment and control strategy based on globally accepted principles and locally applicable strategies are to be followed. These strategies should focus on preventing the development of infection rather than treating diseased stocks.

USE OF ANTIMICROBIALS IN AQUACULTURE

Antimicrobials are used globally both for humans and animals to prevent and treat contagious diseases (O'Neill, 2014). Furthermore, in some countries, antimicrobials are used in animal breeding as growth promoters (Flores et al., 2017). Low and sub therapeutic antimicrobial dosage plays a very important role in improving feeding proficiency, stimulating animal growth, disease avoidance, and control (Magouras et al., 2017). Overall, antibiotic use in the livestock sector is increasing and estimates of total use range from around 63, 000 tons, to over 240, 000 tons per year (Van Boeckel et al. 2015). Antimicrobials are widely employed in aquaculture industry, which mainly contribute to the rise of antimicrobial resistance, sustain potential consequences for animal, human, and ecosystem health. (Daniel Schar et al., 2020) The faster growth of aquaculture has emerged in a series of developments harmful to the human and aquatic animal health as well as environment. Also, the exploit of wide variety human non-biodegradable antibiotics remains aquatic environment. (M.G. Rasul1 and B.C. Majumdar., 2017) It directly causes development of antibiotic-resistance bacteria and transferable resistance genes can be transferred to disease-causing bacteria, resulting in antibiotic-resistant infections for humans, fish and other aquatic animal. The greater the volume of antibiotics used, the greater the risks that antibiotic-resistant populations of

bacteria will induce in the contest for survival of the fittest at the bacterial level. Judicious use of antibiotics in animals is a requirement to delay the emergence of bacteria resistant to the still-working antibiotics, (Vangelis Economou and Panagiota Gousia., 2015) Furthermore, fish farmers must confirm that fish are kept in the best state of health and welfare. The invention of novel drugs or the use of alternatives to antibiotics should also be encouraged. Vast assemblage of antibiotics are applied to the aquaculture in a belief that it will enhance the production and also will improve the socio economic profile in developing countries. (R.Vignesh et al., 2011). It is well acknowledged that the problems relating to antimicrobial use in animal food and aquaculture are of global concern. Intensive fish farming has promoted the outspread of several bacterial diseases, which in turn has led to the abundant use of antimicrobials. The high proportions of antibiotic-resistant bacteria that persevere in sediments and farm environments may provide a threat to fish farms because they can serve as sources of antibiotic-resistance genes for fish pathogens in the vicinity of the farms (Miriam Reverter et al., 2020). In addition to the potential effects on human health, inefficiencies in antibiotic treatment of fish illnesses lead to significant economic losses. One strategy for reducing antibiotic use in aquaculture is to execute rearing practices that minimize the level of stress on the fish and that reduce the likelihood that infections requiring antibiotic treatment will occur. (Ayalew Assefa and Fufa Abunna., 2018).

EVIDENCE OF ANTIMICROBIAL RESIDUES IN AQUACULTURE

Residues of antimicrobials in food have received much surveillance in recent years because of growing food safety and public health concerns. (Fritz Michael Treiber and Heide Beranek-Knauer., 2021). Antimicrobial existence in foods of animal origin constitutes socioeconomic challenges in international trades in animal and animal products. The major public health significances of antimicrobial residues include the development of antimicrobial drug resistance, hypersensitivity reaction, carcinogenicity, mutagenicity, teratogenicity, bone marrow depression, and disruption of normal intestinal flora. Indiscriminate use of antimicrobials in aquaculture resulting in matter of residues in aquaculture products and associated harmful health effects in humans requires control measures to ensure consumer protection. (Reuben Chukwuka Okocha¹ et al., 2018) Ungoverned use of antibiotics in aquaculture industry for the production of farm raised fish and shrimps could pose human health and food safety concerns that remain largely nonregulated in most developing nations of the world. The presence of antibiotic residues in aquaculture products could result in the development of bacterial resistance and toxicity to consumers that can lead to morbidity and or death. Chloramphenicol residues, for example, lead to an increased risk of developing cancer and in very low concentrations may trigger aplastic anemia, a disease that causes bone marrow to stop producing red and white blood cells and is often irreversible and fatal. Other toxic effects include immunopathological effects and carcinogenicity by sulphamethazine, oxytetracycline, and furazolidone; mutagenicity and nephropathy by gentamicin; and allergy by penicillin. The presence of antibiotic residues in domestic animal products and the associated consumer's health hazards have been reported with little attention focused on the aquaculture industry. Today antimicrobials play a major

role in modern livestock production for avoidance and treatment of diseases as well as growth fostering (Feiyang Ma et al., 2021). The global rise in production and demand for aquaculture products has resulted in increasing vulnerability on antibiotics with resultant residues in the products produced for human consumption. (Daniel Schar et al., 2020). Antimicrobial residues irrespective of the route or purpose of administration, antimicrobials can accumulate as residues in tissues, before they are completely metabolized or excreted from the body. The occurrence of residues in fish or other animal tissues is most likely when animals are harvested for human utilization. Consumption of such products may result in many health problems in humans (Reuben Chukwuka Okocha et al., 2018). The problem of antimicrobial residues in animal products is not new. However, due to the globalization of the food trade we are constantly fronting new challenges. Educational work regarding antimicrobial residues in animal products and the associated blooming of multi-resistant germs must be enhanced in developing countries. (Fritz Michael Treiber and Heide Beranek Knauer., 2021). This can only be reduced through increased monitoring of the use of antibiotics in animal breeding. In any case, users must be informed about the health consequences for their customers regarding antibiotic residues in animal products and the development of multi-resistant germs. While low level residues of certain antibiotics are considered safe in some food products, residues of other antibiotics may pose inappropriate risk to public health and are therefore forbid for use in food animals. (Ian Philips et al., 2004). Without antibiotics, food animals suffering from bacterial infectious diseases will be denied effective treatment and outbreaks of disease may not be effectively controlled or prevented within a herd. (Christy Manyi et al., 2018). In general, antibiotics should only be used when preferable and used in food animals; they should be under veterinary supervision. Aquaculture products should be screened for safety levels of antimicrobial residues. Efforts should also be made to reduce the use of antibiotics by implementing good animal husbandry practices.

PUBLIC HEALTH IMPACT OF ANTIMICROBIALS

Aquaculture has become the fastest growing sector of food production in the world. Despite the encouraging trends, several obstructions have negative impact on the growth of aquaculture. Therefore, strict measures, legislations and regulations for the use of antimicrobials in aquaculture should be developed and implemented, especially in developing countries, to avoid such negative impacts in human, fish, animals and environment. (Salah Mesalhy Aly and Aqel Albutti., 2014). The intensification of aquaculture to achieve market demands could lead to an increase in infectious diseases by pathogenic bacteria. Consequently, antimicrobials act as controls for emerging infectious diseases, but their use must follow the rules and regulations of the country where the activity is performed. (Louis Eduardo de Souza Gazal et al., 2020). Resistant bacteria from fish farming are a serious concern because they can be acquired by humans with handling or food chain, which may represent a public health problem. Excessive use of antimicrobials in aquaculture can thus potentially negatively impact animal and human health as well as the aquatic environment and should be better evaluated and regulated. (Felipe C Cabello et al., 2013). Antibiotics have been

extensively and effectively used in aquaculture due to expedite growth of aquaculture and this has resulted in the development of serious health problems in aquaculture, other animals, and human. (Pathmalal M Manage., 2018).

ALTERNATE ANTIBIOTIC STRATEGIES IN AQUACULTURE

The wide and frequent use of antibiotics in aquaculture has resulted in the development and spread of antibiotic resistance. Because of the health risks associated with the use of antibiotics in animal production, there is a growing awareness that antibiotics should be used with more care. This is reflected in the recent execution of more strict regulations on the prophylactic use of antibiotics and the presence of antibiotic residues in aquaculture products. (Tom Defoirdt et al., 2011). The aquatic environment is more supportive to pathogenic bacteria independently of their host than the terrestrial environment and pathogens can reach high densities around the animals, which then ingest them either with the feed or when they are drinking. Culturing several species of aquatic animals in many cases suffers from highly uncertain survival rates because of bacterial diseases (Laurent Verschuere et al., 2000). In addition to regulations on the use of antibiotics in aquaculture, many governments around the world have enforced Maximum Residue Levels (MRLs) for aquaculture products (Ronald Lulijwa et al., 2019). One of the alternative strategies that has recently been developed to control infections caused by antibiotic-resistant bacteria is the disruption of quorum sensing, bacterial cell-to-cell communication. (Tom Defoirdt et al., 2007). This is in contrast to conventional antibiotics, which also affect harmless and beneficial bacteria and pose strong selective pressure in any environment. Indeed, the biocontrol agents include viruses (bacteriophages), algae or their metabolites and bacteria or their metabolites. In this respect, the alternative strategies discussed here could be the basis for a rational selection of probiotics bacteria based on different modes of action. Indeed, selection of probiotics for aquaculture so far focused mainly on the production of constraint substances in vitro. (S. Mohapatra et al., 2012). Although most of them have been tested both in vitro and in vivo, few have been tested in ‘real life’ situations. Because most probably no biocontrol strategy will be successful in all cases, we think that it is important to further develop different strategies that could be combined or used in rotation in order to maximize the chance of successfully protecting the animals and to prevent resistance development.

CURRENT STATUS AND FUTURE CHALLENGES IN AQUACULTURE

The results obtained by using techniques that disrupt quorum sensing systems of pathogenic bacteria indicate that it is a promising replacement for antibiotics in fighting bacterial infections. (Tom Defoirdt et al., 2004). This new approach might also have value in aquaculture since a link between quorum sensing and virulence factor expression in several aquatic pathogens has been demonstrated. Unfortunately, data about the impact of quorum sensing on virulence (i.e. the net result of all virulence factors) of aquatic pathogens are still lacking. Hence, bacterial diseases affecting either agricultural crops, fish, or shellfish not only cause large economic losses to producers

but can even create food shortages, resulting in malnutrition, or even famine, in vulnerable populations. Phages are viruses that specifically infect bacteria; they are highly host-specific and represent an environmentally-friendly alternative to antibiotics to control and kill pathogenic bacteria.

CONCLUSION

The diverse nature of aquaculture leads to the paradox that it is often proposed as a solution to global food supply issues, but may actually be a net consumer of fish and reduce the availability of fish for people (Naylor et al., 2000) culture where predators and competitors are controlled. Although the boom that took place in the aquaculture sector in the eighties and early nineties, crashed in the mid-nineties but still an opportunity exists for revival of the industry. What is required is the practice of responsible and technologically sound aquaculture that will be eco-friendly and sustainable. The significant growth in fisheries and aquaculture production, especially in the past two decades, has enhanced the world's capacity to consume diverse and nutritious food. Fish and fish products have a crucial role in nutrition and global food security, as they represent a valuable source of nutrients and micronutrients important for diversified and healthy diets. Moreover, in lower-income countries, the importance of fish is enhanced by the fact that it contains many of the vitamins and minerals required to address some of the most severe and widespread nutritional deficiencies. Even small quantities of fish in people's diet can have a significant positive impact on the nutritional status of poor consumers across the world. Since 1961, the annual global growth in fish consumption has been twice as high as population growth, demonstrating that the fisheries sector is crucial for a world without hunger and malnutrition (FAO, 2018). Despite the significant contributions that fisheries and aquaculture make to employment, nutrition, and trade in the developing world, they are rarely included in national development policy and donor priorities. This is largely due to problems with valuation of small-scale fisheries, as policy makers often do not have access to data which reflect the importance of fisheries and aquaculture to development. Even with improvements in regulation, however, pressures on capture fisheries will remain, due to continued population growth. Further development of sustainable aquaculture and improvements in the post-harvest sector to reduce losses could help to maintain fish supply and the contribution of fish.

References

- [1] Roslina Kamaruddin and Amir Hussin Baharuddin (2015), "The importance of good aquaculture practices in improving fish farmer's income" A case of Malaysia, Department of Economics and Agribusiness, Article in International Journal of Social Economics December 2015.
- [2] Michael Tlusty Edgerton (2002), "The benefits and risks of aquacultural production for the aquarium trade Research Laboratory" New England Aquarium, Central Wharf, USA Aquaculture 205 (2002) 203–219.

- [3] M.Kumaran, V.S.Chandrasekaran and N. Kalaimani (2003), "Aquaculture Extension :the neglected aspect in Aquaculture Development" Central Institute of Brackishwater Aquaculture (CIBA), Santhome High Road, Chennai Fishing Chimes Vol.23 (2003).
- [4] Sudipto Sarkar and Bimal Chandra Mal (2005), "The Status of Aquaculture in India : Development, Adoption and Constraints" Agricultural Engineering Today, Vol. 29 (5-6): 46-52, 2005
- [5] M. Krishnan and Pratap S.Birthal (2002), "Aquaculture development in India: an economic overview with special reference to coastal aquaculture Central Institute of Brackishwater Aquaculture", Chennai, India Aquaculture Economics and Management 6 (1/2) 2002
- [6] P. Jayasankar, K.N. Mohanta and S. Ferosekhan (2018), "Freshwater Aquaculture in India", ICAR Central Institute of Freshwater Aquaculture, Odisha Aquaculture in India, Pages 23-49 2018.
- [7] MD. Ashikur Rahman, Shirin Akter, M Mominuzzaman Khan and MD.Khalilur Rahman (2019), "Relation between aquaculture with fish disease & health management: A review note", Bangladesh Fisheries Research Institute, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj Bangladesh J. Fish. (2019) 31 (2) : 253-260.
- [8] Ayalew Assefa and Fufa Abunna Rahman (2018), "Maintenance of Fish Health in Aquaculture: Review of Epidemiological Approaches for Prevention and Control of Infectious Disease of Fish", Sekota Dryland Agricultural Research Center, Ethiopia College of Veterinary Medicine and Agriculture, Oromia, Ethiopia Veterinary Medicine International Volume 2018, Article ID 5432497, 10 pages.
- [9] Daniel Schar1, EiliY. Klein, Ramanan Laxminarayan, MariusGilbert and Thomas P.Van Boeck (2020), "Global trends in antimicrobial use in aquaculture", Spatial Epidemiology Laboratory, Belgium, Center for Disease Dynamics, Washington, USA, Institute for Environmental Decisions, Switzerland.Scientific Reports | (2020) 10:21878.
- [10] Esther Eyram Agoba, Francis Adu, Christian Agyare and Vivian Etsiapa Boamah Boeck (2017), "Antibiotic Use and Practices in Selected Fish Farms in the Ashanti Region of Ghana", Department of Pharmaceutics, Faculty of Pharmacy and Pharmaceutical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana J Infec Dis Treat. 2017, Vol.3 No.2:9.
- [11] Jaime Romero, Carmen Gloria Feijoó and Paola Navarrete (2012), "Antibiotics in Aquaculture—Use, Abuse and Alternatives", Universidad de Chile, Biotechnology Laboratory, Institute for Nutrition and Food Technology (INTA) Universidad Andrés Bello, Laboratory of Developmental Biology, Santiago Chile Health and Environment in Aquaculture. 2012
- [12] Dr. Charles M. Benbrook, (2002), "Antibiotic Drug Use in U.S. Aquaculture", The Northwest Science and Environmental Policy Center Sandpoint, Idaho Antibiotic Use in Aquaculture. 20 13. M.G. Rasul1, B.C. Majumdar Dr. Charles M. Benbrook, (2017), "Abuse of Antibiotics in Aquaculture and it's Effects on Human, Aquatic Animal and Environment", Department of Fisheries

- Technology, Bangabandhu, Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh Haya: Saudi J. Life Sci.; Vol-2, Iss-3 (Apr-Jun, 2017):81-88
- [13] Pathmalal M. Manage, (2018), "Heavy use of antibiotics in aquaculture:Emerging human and animal health problems–A review", Centre for Water Quality and Algae Research, Department of Zoology, University of Sri Jayewardenepura, Nugegoda, Sri Lanka, J. Aquat. Sci. 23 (1) (2018): 13-27
- [14] R. Vignesh, B.S. Karthikeyanb, N. Periyasamy and K. Devanathan, (2011), "Antibiotics in aquaculture: An overview", Centre for Advanced Study in Marine Biology, Faculty of Marine Sciences, Annamalai University, Tamilnadu, India, Department of Bioinformatics, School of Life Sciences, Bharathidasan University, Tamilnadu, India, J Exp Biol; 1 (3): 2011
- [15] Felipe C. Cabello, (2006), "Heavy use of prophylactic antibiotics in aquaculture:a growing problem for human and animal health and for the environment", Department of Microbiology and Immunology, New York Medical College, USA, Environmental Microbiology (2006) 8 (7), 1137–1144
- [16] Marta Sargenti, Silvia Bartolacci, Aurora Luciani, Katiuscia Di Biagio, Marco Baldini, Roberta Galarini, Danilo Giusepponi and Marinella Capuccella, (2020), "Investigation of the Correlation between the Use of Antibiotics in Aquaculture Systems and Their Detection in Aquatic Environments: A Case Study of the Nera River Aquafarms in Italy", Experimental Zooprophylactic Institute of Umbria and Marche, Italy Sustainability 2020.
- [17] Fritz Michael Treiber and Heide Beranek-Knauer, (2021), "Antimicrobial Residues in Food from Animal Origin A Review of the Literature Focusing on Products Collected in Stores and Markets Worldwide", Institute of Molecular Biosciences, University of Graz, Austria Antibiotics 2021, 10, 534.
- [18] Reuben Chukwuka Okocha¹, Isaac Olufemi Olatoye and Olufemi Bolarinwa Adedeji, (2018), "Food safety impacts of antimicrobial use and their residues in aquaculture", Department of Veterinary Public Health and Preventive Medicine, University of Ibadan, Nigeria, Public Health Reviews (2018) 39:21.
- [19] G.K.Sivaraman V, Murugadas and M.M.Prasad (2018), "Impact of Antimicrobial resistance (AMR) in aquatic products", Microbiology, Fermentation and Biotechnology Division, ICAR-Central Institute of Fisheries Technology, Cochin–Kerala. ICAR-CIFT, Cochin, 05-25, 2018.
- [20] Anwesha Roy, Avijit Biswas, Subal Kumar Ghosh, Hauzou Kim and Sagarika Swain (2021), "Current scenario of antibiotic-resistance in Indian aquaculture", School of Fisheries, Centurion University of Technology and Management, Odisha, India, The Pharma Innovation Journal 2021; SP-10 (7): 454-461
- [21] Lucia Santos, Fernando Ramos (2018), "Antimicrobial resistance in aquaculture:Current knowledge and alternatives to tackle the problem", Pharmacy Faculty, University of Coimbra, Portugal International Journal of Antimicrobial Agents 52 (2018) 135–143
- [22] Lucia Santos, Fernando Ramos (2018), "Antimicrobial resistance in aquaculture:Current knowledge and alternatives to tackle the problem",

- Pharmacy Faculty, University of Coimbra, Portugal *International Journal of Antimicrobial Agents* 52 (2018) 135–143
- [23] Ole E. Heuer, Hilde Kruse, Kari Grave, P. Collignon, Iddya Karunasagar and Frederick J. Angulo (2009),), “Human Health Consequences of Use of Antimicrobial Agents in Aquaculture”, Danish Zoonosis Centre, National Food Institute, Denmark; National Veterinary Institute and Department of Food Safety and Infection Biology, Norwegian School of Veterinary Science, Norway *FOOD SAFETY CID* 2009.
- [24] Carmen Sieiro, Lara Areal-Hermida, Ángeles Pichardo-Gallard, Trinidad de Miguel, Sandra Sánchez, Ángeles Sánchez-Pérez and Tomás G (2020),), “A Hundred Years of Bacteriophages: Can Phages Replace Antibiotics in Agriculture and Aquaculture?”, Department of Functional Biology and Health Sciences, Spain, Sydney School of Veterinary Science, Faculty of Science, University of Sydney, Australia, *Antibiotics* 2020.
- [25] Pragyan Dash, Satheesha Avunje, Ritesh S. Tandel, Sandeep K. P. and Akshaya Panigrahi (2017), “Biocontrol of Luminous Vibriosis in Shrimp Aquaculture: A Review of Current Approaches and Future Perspectives”, Indian Council of Agricultural Research-Directorate of Coldwater Fisheries Research (DCFR), Bhimtal, India; Indian Council of Agricultural Research-Central Institute of Brackishwater Aquaculture (CIBA), Chennai, India, *Reviews in Fisheries Sciences and Aquaculture* 2017.
- [26] Relation between aquaculture with fish disease & health management: A review note MD. Ashikur Rahman, Shirin Akter, M Mominuzzaman Khan and MD.Khalilur Rahman Fisheries Research Institute, Freshwater Station, Mymensingh, Bangladesh Department of Fisheries and Marine Bioscience, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj J. Fish. (2019) 31 (2) : 253-260
- [27] Present Status of Fish Disease Management in Freshwater Aquaculture in India Mishra, Rakesh D, Dhiman M, Choudhary P, Debbarma J, Sahoo SN, Barua A, Giri Ramesh R, Ananda K, Mishra CK and Swain P Fish Health Management Division, ICAR-Central Institute of Freshwater Aquaculture (CIFA), Kausalyaganga, Bhubaneswar, Odisha, India *Journal of Aquaculture & Fisheries* 2017
- [28] Maintenance of Fish Health in Aquaculture: Review of Epidemiological Approaches for Prevention and Control of Infectious Disease of Fish Ayalew Assefa and Fufa Abunna Dryland Agricultural Research Center, Sekota, Ethiopia, *Veterinary Medicine International*, vol. 2018.
- [29] More rapid and severe disease outbreaks for aquaculture at the tropics: implications for food security Tommy L. F. Leung and Amanda E. Bates Centre for Behavioural and Physiological Ecology, Zoology, University of New England, Australia ; *Journal of Applied Ecology* 2013
- [30] Global trends in antimicrobial use in aquaculture Daniel Schar, EiliY. Klein, Ramanan Laxminarayan, MariusGilbert1&Thomas P.Van Boeckel, *Scientific report*, Nature 2020

- [31] Antibiotics in Aquaculture—Use, Abuse and Alternatives Jaime Romero¹, Carmen Gloria Feijoó and Paola Navarrete Universidad de Chile, Biotechnology Laboratory, Institute for Nutrition and Food Technology (INTA) Universidad Andrés Bello, Laboratory of Developmental Biology, Santiago Chile Health and Environment in Aquaculture 2012
- [32] Abuse of Antibiotics in Aquaculture and it's Effects on Human, Aquatic Animal and Environment M.G. Rasul¹, B.C. Majumdar¹ ¹Department of Fisheries Technology, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh ; Haya: Saudi J. Life Sci.; Vol-2, Iss-3 (Apr-Jun, 2017):81-88
- [33] Unravelling the menace: detection of antimicrobial resistance in aquaculture P.G. Preena, T. Raja Swaminathan, V.J. Rejish Kumar and I.S. Bright Singh National Centre for Aquatic Animal Health, Cochin University of Science and Technology, Kochi, India Lett Appl Microbiol 2020.
- [34] Evaluating antimicrobial resistance in the global shrimp industry Kelly Thornber, David Verner-Jeffreys, Steve Hinchliffe¹ Muhammad Meezanur Rahman, David Bass and Charles R. Tyler Centre for Sustainable Aquaculture Futures, University of Exeter, UK, Worldfish Bangladesh, World Fish Bangladesh Office, Dhaka, Bangladesh, Rev Aquac 2020.
- [35] The Rising Tide of Antimicrobial Resistance in Aquaculture Joy E. M. Watts, Harold J. Schreier, Lauma Lanska and Michelle S. Hale School of Biological Sciences, University of Portsmouth, UK; Departments of Marine Biotechnology and Biological Sciences, University of Maryland Baltimore County, Baltimore, USA 2017
- [36] Aquaculture at the crossroads of global warming and antimicrobial resistance Miriam Reverter, Samira Sarter Domenico Caruso, Jean-Christophe Avarre, Marine Combe, Elodie Pepey, Laurent Pouyaud, Sarahi Vega-Heredía Hugues de Verdal Rodolphe E. Gozlan April 2020 Nature Communications.
- [37] Raising awareness of antimicrobial resistance in rural aquaculture practice in Bangladesh through digital communications: a pilot study Kelly Thornber, Doina Huso, Muhammad Meezanur Rahman, Himangsu Biswas, Mohammad Habibur Rahman, Eric Brum & Charles R. Tyler 2019
- [38] Antimicrobial resistance in aquaculture: a crisis for concern Prasannan Geetha Preena¹ & Thangaraj Raja Swaminathan¹ & Vattiringal Jayadrathan Rejish Kumar² & Isaac Sarojini Bright Singh 2019
- [39] Use of molecular and genomic data for disease surveillance in aquaculture: Towards improved evidence for decision making. Katharina D.C. Stärk, Agnieszka Pękala, Petra Muellner Switzerland Royal Veterinary College, London, UK National Veterinary Research Institute, Puławy, Poland. Prev Vet Med 2019
- [40] Heavy use of antibiotics in aquaculture: Emerging human and animal health problems Pathmalal M. Manage Centre for Water Quality and Algae Research, Department of Zoology, University of Sri Jayewardenepura, Nugegoda, Sri Lanka J. Aquat. Sci. 2018)

- [41] Antimicrobials and resistant bacteria in global fish farming and the possible risk for public health Eduardo de Souza Gazal Kelly Cristina Tagliari de Brito², Renata Katsuko Takayama Kobayashi¹, Gerson Nakazato, Lissandra Souto Cavalli³, Luciana Kazue Otutumi, Benito Guimarães de Brito *Arq. Inst. Biol.*, v.87, 1-11, e0362019, 2020
- [42] Fresh water Aquaculture in India P. Jayasankar, K.N. Mohanta and S. Ferosekhan ICAR Central Institute of Freshwater Aquaculture, Kausalyganga, Bhubaneswar, Odisha Aquaculture in India 2018
- [43] Antibiotic use and practices in Selected Fish Farms in the Ashanti Region of Ghana Esther Eyram Agoba, Francis Adu, Christian Agyare and Vivian Etsiapa Boamah Antibiotic Use and Practices in Selected Fish Farms in the Ashanti Region of Ghana. *J Infec Dis Treat.* 2017.
- [44] Antibiotics in Aquaculture—Use, Abuse and Alternatives Jaime Romer, Carmen Gloria Feijo and Paola Navarrete Universidad de Chile, Biotechnology Laboratory, Laboratory of Developmental Biology, Santiago Chile Health and Environment in Aquaculture 2012
- [45] Antibiotic Drug Use in U.S. Aquaculture Dr. Charles M. Benbrook The Northwest Science and Environmental Policy Center Sandpoint, Idaho February 2002
- [46] Current scenario of antibiotic-resistance in Indian aquaculture Anwesha Roy, Avijit Biswas, Subal Kumar Ghosh, Hauzou Kim and Sagarika Swain *The Pharma Innovation Journal* 2021
- [47] A 20-year retrospective review of global aquaculture Rosamond L. Naylor, Ronald W. Hardy, Alejandro H. Buschma, Simon R. Bush, Ling Cao, Dane H. Klinger, David C. Little, Jane Lubchenco, Sandra E. Shumway & Max Troell, *Nature* 2021 Mar
- [48] The Rise and Fall of Antibiotics in Aquaculture Antony T. Vincent, Jeff Gauthier, Nicolas Derome, and Steve J. Charette *Microbial Communities in Aquaculture Ecosystems, Improving Productivity and Sustainability* 2019