

Effect of geomagnetic storms and their association with solar wind velocity and IMF during solar cycle 23 and 24

Sham Singh^{*1}, A. C. Panday², Kalpana Singh³ & A. P. Mishra²

¹*Department of Applied Sciences, Chandigarh Engineering College,
Landran, Mohali, Punjab, India.*

²*Department of Physics, A. P. S. University, Rewa -486003 (M.P), India.*

³*Department of Applied Sciences & Humanities, Kamla Nehru Institute of Technology,
Sultanpur (U.P.), India.*

Abstract

In this paper we present the results on interplanetary causes of large geomagnetic storms ($Dst \leq -100nT$ to $Dst \leq -200nT$), that occurred during solar cycle 23 and 24 (1996–2012). It was reported that the most common interplanetary structures leading to the development of large storms were: magnetic clouds, sheath fields, sheath fields followed by a CMEs and solar flares regions at the leading fronts of high speed streams. However, the relative importance of each of those driving structures has been shown to vary with the solar cycle phase. Space weather in the near-earth environment is directly associated to disturbances originating at the Sun. Two interplanetary structures are important for the development of storms, involving intense southward IMFs: the sheath region just behind the forward shock, and the CME eject itself, around solar maximum. During solar minimum, the high-density, low-speed streams associated with the heliospheric current sheet (HCS) plasma impinging upon the Earth's magnetosphere cause positive Dst Values.

Keywords: Solar wind velocity, Average of Interplanetary Magnetic field IMF (B), CME's and geomagnetic storms.

INTRODUCTION

Strong interaction of CMEs with Earth's environment causes serious space weather effect through the coupled magnetosphere system. A geomagnetic storm is a global disturbance of the earth's magnetic field (Akasofu, 1963), and usually occurs in response to abnormal conditions in the IMF and solar wind. The geomagnetic storms are a sequence of varying magnetospheric response to the varying conditions in interplanetary space which are caused by coronal magnetic storms. A variety of geomagnetic storms and several sources of their origin have been suggested by many authors (Tsurutani et al, 1988). The phenomenon of the geomagnetic storm and its manifestation has been discussed in the literature (Chapman 1936; Murayama 1982; Oh and Yu 2004; Jatin Rathod et al. 2008 and Kane 2014; Singh and Mishra 2015). Hewish and Bravo (1986) found that geomagnetic storms are more associated with coronal holes than the solar flares. Burlaga et al. (1987) have concluded that these are associated with either a magnetic cloud which is also confirmed by Wilson (1987).

The impacts of an interplanetary irregularity causing a sudden change in the geomagnetic components are called sudden commencements (Smith et al. 1986). It is accepted that the impact of the interplanetary irregularity on the magnetopause causes hydromagnetic waves which travel through the magnetosphere towards the Earth as also into the tail and their isotropic modes give the SSC. The importance of studying geomagnetic storms is basically two fold. One refers to their academic aspect and the other involves principal aspects that in some cases can represent a particular concern for mankind. The study of geomagnetic storms is useful to explore adverse effect in radio communications, power grid, radar observations, electrical utilities, long-distance pipelines and synchronous spacecraft. Geomagnetic activity can be divided in to two main categories, storms and sub storms. Storms, the main contributors to space weather, are initiated when enhanced energy transfer from the solar wind/ IMF to the magnetosphere leads in to intensification of ring current. The ring current development can be monitored with the Dst index (Gonzales et al., 1994; Bakare 2010; Kane 2014).

Some of the most dramatic space weather effects occur in association with eruption of material from the solar atmosphere into interplanetary space. These eruptions are known as coronal mass ejections, (CMEs). A large CME can contain 10^{16} grams (a billion tons) of matter that can be accelerated to several million miles per hour in a spectacular explosion. CMEs originating from close to the disk center significantly perturb earth's environment and they directly impact the earth (Gopalswamy 2006). Each CMEs are drain the solar mass in the range of 10^{10} Kg (Webb. 1995; Singh et al. 2012). Kahler (1992) pointed out if the CME is associated with flare then the CME originates in the explosive phase of the flare. Gosling (1993) showed that solar flares play no fundamental role in causing geomagnetic disturbances. Flares and CMEs are part of the same magnetic eruption process (Harrison 1995; Schreiber 1998; Zhang et al. 2001; Singh et al. 2012; Singh et al 2013). Characteristics of flares associated with CMEs has been discussed in the literature (Mishra et al. 2008; Balveer 2011; Singh and Mishra 2015).

SELECTION CRITERIA AND DATA ANALYSIS

In the present study, we have analyzed the Geomagnetic storm with Dst magnitude ≤ -100 nT and their relationship with solar flare, and coronal mass ejections, occurred during solar cycle 21 to 23. The daily observed H- α solar flare data has been taken from the GOES Satellite which is supported by National Geophysical Data Center (NGDC). CMEs are routinely observed by the LASCO telescope on board the SOHO satellite. The extrapolated CMEs onset time is close to the associated type II and IV bursts. The properties of CMEs are collected from the data available through website; <http://www.cdaw.gsfc.gov/cmelist/>. The catalog contains a list of all visible CMEs with information of their date and time of the first appearance in the field of view of c₂ coronagraph, central position angle, angular width, speed acceleration obtained from quadratic fitting etc, and the Dst, solar wind, IMF, Bz data has been taken from the omni web data.

RESULTS AND DISCUSSION

The study has once again confirmed the earlier results (Gonzalez et al., 1994, Dubey and Mishra 2000; Singh and Mishra 2015) that the indices which are associated with long term variation of sunspot activity sometimes show different behaviour on short term scale depending on local solar active regions and associated phenomena. Since these individual indices on a day to day basis, are affected differently at different times, hence we have tried to study the major geomagnetic storms as an individual event associated with solar and interplanetary disturbances. Here we have selected only those geomagnetic storms whose Dst magnitude ≤ -100 nT, occurred during the last three decades.

The largest interplanetary disturbances, in particular the coronal mass ejections (CME) and the high-speed solar wind streams (HSS) are known to cause dramatic effects in the near-Earth space, accelerating magnetospheric (in particular, auroral and ring current) particles to higher energies and enhancing particle precipitation into the atmosphere, especially in the mesosphere-upper thermosphere (MLT) region. Moreover, magnetospheric particle precipitation maintains and produces electric currents and fields that also affect the charged and neutral atmosphere below. The effects are found to depend on the nature of the interplanetary disturbance and to be very different for CMEs and HSSs.

The geomagnetic storms preceded by sudden commencement have been divided into three phases, the initial phase, main phase and recovery phase. The initial phase is caused by an enhancement of solar wind behind the shock wave. It is quasi steady state preceded by sudden storm commencement. The main phases of geomagnetic storms follow with the sudden ionospheric disturbances. The recovery phases of geomagnetic storm follow with the active main phase. It consists of slow quite return of H field back to pre storm level. It is also noted that recovery phase duration is always higher than the main phase duration. In **Fig. 1** shows the total number of large geomagnetic storms observed during said period is 88, associated CMEs with

geomagnetic storms is 43 and associated CMEs + flare are 18 and associated with only flare is 27.




Figure 1: Shows the large Geomagnetic Storms and their association with CMEs, CMEs + Flares and Flares.

GEOMAGNETIC STORM OF 15 DECEMBER 2006

To obtain further insight on the effectiveness of various parameters, we have analysed the nature of effects during various events through an examination of the profiles of all the interplanetary variables. Earlier many workers have studied the major geomagnetic storms and their association with different solar & interplanetary features (Singh and Mishra, 2015). Solar and interplanetary disturbances causing moderate geomagnetic storms (Santosh Kumar et al. 2008). The events have been distributed according to the magnitude and nature of the profile in each case. The changes in these parameters have been grouped in different slots of magnitudes.

During the geomagnetic storm, the solar wind velocity has an increasing tendency with change of velocity in the range up to 550 to 650 km/sec (56 cases) which in some cases rises to about 650 – 750 km/sec and rarely sometimes \approx 900 km/sec during the event of 15 Dec. 2006 **Fig.2**. Whether these CMEs were actually frontside events but lacked visible surface signatures, were backside events with global consequences such as expulsions triggered by global waves (Mc Comas et al. 1991).




Figure 2: Shows the variation of solar wind velocity during 12 Dec. to 18 Dec. in year 2006.

There is also abrupt change in Bz component from northward to southward. Not only negative Bz or B average is responsible for development of MGS but also its magnitude and duration contribute significantly (Kane 1977). **Fig. 3** shows the variation in Bz component during the some event. The southward Bz component becomes prominently negative indicating the geomagnetic storm occurrence.




Figure 3: Shows the variation of Bz components of IMF during 12 Dec. to 18 Dec. in year 2006.

The variability of Dst parameters for each day is given in figure (4) during geomagnetic storms 15 December 2006 (Dst peak magnitude -140 nT). In phase with the ring current excursion, however, Murayama (1982) found a correlation between negative Dst and dynamic pressure, which was recently confirmed by Fenrich and Luhmann (1998), but this dependence of storm strength on dynamic pressure is weak. Peak magnitude occurred at 7 UT on 15 Dec. 2006, and recovered to normal level on 18 Dec. 2006. At last this geomagnetic storm has been found to be associated with the 4B flare / CME (Speed 1042 km/sec.) This event has occurred during the declining phase of solar cycle 23, which is a remarkable feature.




Figure 4: Shows the geomagnetic storms during the time period of 12 Dec. to 18 Dec. in year 2006.

From **fig. 5** and **6** it is apparent that total magnetic field (B) has good correlation with Kp and Ap Indices Correlation Coefficient $r = (0.59)$ and (0.58) respectively.




Figure 5: Correlation plots of geomagnetic index Ap with total average interplanetary magnetic IMF (B) field during the period 12-18 April, 2006.




Figure 6: Correlation plots of geomagnetic index Kp with total average interplanetary magnetic field IMF (B) during the period 12-18 April, 2006.

CONCLUSION

On the basis of results and discussion, we have summarized important results derived from the analysis of the geomagnetic storms (of magnitude <-100 nT) and their relationship with interplanetary parameters for the period 1996 to 2012 as follows:

It is found that total number of large geomagnetic storms ($Dst \leq -100$ nT) were observed 88. WE conclude that large storms associated with CMEs were 43, associated with CMEs and solar flares (B-class and M-class), and however associated with only flare is 27.

Major Geomagnetic storms observed on 14 April 2006 have been studied and their relationship with interplanetary parameters has also been discussed. The total average interplanetary magnetic field (B) has good correlation with Kp and Ap Indices with Correlation Coefficient $r = (0.59)$ and (0.58) respectively. These quantitative relationships are invaluable for modeling studies and space weather phenomena. It has been verified that geomagnetic storm intensity is correlated well with the total magnetic field of IMF better than southward component B_z of the IMF, density and solar wind velocity.

REFERENCES

- [1] Akasofu, S.I. and Chapman, S. (1963) *Space Sc. Rev.* 2, 91.
- [2] Balveer S. Rathore, Subhash C. Kaushik, K. A. Firoz, D. C. Gupta, A. K. Shrivastava, Krishna Kant Parashar, and Ram Mohan Bhaduriya. (2011) A Correlative Study of Geomagnetic Storms Associated with Solar Wind and

- IMF Features During Solar Cycle 23. *International J. of Applied Phys. and Mathematics*, 1, 2, 149-154.
- [3] Bakare, N. O. and Chukwuma, V. U. (2010) Relationship between Dst and solar wind conditions during intense geomagnetic storms. *Indian J Radio Space Phys*, 39, 150-155.
- [4] Biswas, S., Shome, A., Raha, B., Bhattacharya, A.B. (2016) *Indian J. Phys.* 1-8, doi:10.1007/s12648-016-0893-y.
- [5] Burlaga, L.F., Behannon, K.W., Klien, L.W. (1987) *J. Geophys. Res.*, 92, 5725.
- [6] Carlo Benna, Salvatore Mancuso, Silvio Giordano, Lorenzo Gioannini. (2013) Plasma properties from the multi-wavelength analysis of the November 1st 2003 CME/shock event. *J. of Advanced Res.*, 4, 293–296.
- [7] Chapman, S. (1936) *The Earth's Magnetism*, Methuen Monograph on Physical Subjects, Methuen and Co. Ltd., London, U.K.
- [8] Dubey, S.C. and Mishra A.P., (2000) *Indian J. of Radio and Space Phys.*, 29
- [9] Fenrich, F.R., Luhmann, J.G., (1998) *Geophysical Research Letters*. 25, 2999-3002.
- [10] Gopalswamy, N. (2006) *J. Astrophys. Astro.*, 27, 243.
- [11] Gosling, J.T. (1993) *J. Geophys. Res.*, 98(a11), 18, 937-950.
- [12] Gonzalez, W.D., Joselyn, J. A., Kamide, Y., Kroehl, H.W., Rostoker, G., Tsurutani, B.T. and Vasyliunas V.M., 1994, *J. Geophys. Res.*, 99, 5771 – 5792.
- [13] Harrison, R.A. (1995) *A and A*, 304, 585.
- [14] Jatin Ratod, Girija Rajaram, Radharani Alyana, A.Chandrasekhar Reddy, D.S.Misra, C.G. Patil, M.Y.S. Prasad and A.G. Ananth., (2008) *J. Astrophys. Astr*, 29, 293-302.
- [15] Kane, R. P., (2014) Evolution of solar indices during the maximum of solar cycle 24. *Indian J Radio Space Phys*, 43, 151-155.
- [16] Kane, R.P. (1977) *J. Geophys. Res.*, 82, 561.
- [17] Kahler, S.W. (1992) *Annual Review of A and A*, 30 (A93-25826 09-90), 113-141
- [18] Malini Aggrawal, Rajmal Jain, A. P.Mishra, P. G. Kulkarni, Chintan Vyas, R.Shama and Meera Gupta. (2008) *J. Astrophys. Astr*. 29, 195-205.
- [19] Mc Comas, D.J., Phillips, J.L., Hundhausen, A.J., Burkepile, J.T., (1991) *Geophysical Research Letters*, 18, 73-76.
- [20] Murayama, T. (1982) Coupling function between solar wind parameters and geomagnetic indices, *Rev. Geophys. Space Phys.*, 20, 623-639.
- [21] Oh, Yeon, Yu, Yi (2004) Relationships of the solar wind parameters with the magnetic storm magnitude and their association with the interplanetary shock, *J. Korean Astronomical Society*, 37, 151-157.
- [22] Santosh Kumar, M.P. Yadav and Amita Raizada, 2008, *J. Astrophys. Astro*, 29, 263-267.

- [23] Schreiber, H., (1998) On the periodic variations of geomagnetic activity indices Ap and ap. *Ann. Geophysicae*, 16, 510-517.
- [24] Singh, S. and Mishra, A.P. (2015) Interaction of solar plasma near-Earth with reference to geomagnetic storms during maxima of solar cycle 24. *Indian J. Phys.*, 89, 1227-1234.
- [25] Singh, Divya Shrivastava, Shri Ram Lahauriya, Adya Prasad Mishra (2012) Comparative study of solar cycles 22 to 24 in relation to solar output variability. *Natural Science* 4, 6, 349-354. doi:10.4236/ns.2012.46048.
- [26] Singh, S., Shrivastava, D. and Mishra, A. P. (2012) Effect of solar and interplanetary disturbances on space weather. *Indian J. Sci. Res.* 3, 2, 121-125.
- [27] Singh, S., Pandey, A., Singh, K., Mishra, A. P. (2013) Characteristic Features of Geomagnetic Storms Observed During Maxima of Solar Cycle 24. *International Journal of Physics and Astronomy*, 26, 1, 1103-1110.
- [28] Smith, E.J., Slavin, J.A., Zwicki, R.D., Bame, S.J. 1986, Solar wind magnetospheric coupling (eds) Y. Kamide and J.A. Slavin, Terra Scientific. Tokyo, Japan.
- [29] Tsurutani, B. T., Gonzalez, W.D., Tang, F., Akasofu, S. I., Smith E., 1988, *J. Geophys. Res.* 93, 13 851
- [30] Webb, D.F. (1995) *Review of Geophysics*, 33, 577.
- [31] Wilson, Robert M. (1987) *Planet Space Sci.* 35, 339.
- [32] Zhang, J., Dere, K.P., Howard, R.A., Kundu, M.R., White, S. M. (2001) *Astrophys. J.*, 559, 452.

